

2

NPS55-90-17

# NAVAL POSTGRADUATE SCHOOL

Monterey, California

AD-A230 255



DTIC  
ELECTE  
JAN 03 1991  
S E D

MODELING AND SIMULATION OF RANDOM  
SHAPES BY SCULPTURED WRAPAROUND:  
PRELIMINARY REPORT

Donald P. Gaver  
Patricia A. Jacobs

August 1990

Approved for public release; distribution is unlimited.

Prepared for:  
Naval Postgraduate School,  
Monterey, CA 93955

91 1 - 2 034


NAVAL POSTGRADUATE SCHOOL,  
MONTEREY, CALIFORNIA


Rear Admiral R. W. West, Jr.  
Superintendent

Harrison Shull  
Provost

This report was prepared in conjunction with research funded under the  
Naval Postgraduate School Research Council Research Program.


This report was prepared by:


  
DONALD P. GAVER  
Professor of Operations Research

  
PATRICIA A. JACOBS  
Professor of Operations Research

Reviewed by:

Released by:

  
PETER PURDUE  
Professor and Chairman  
Department of Operations Research

  
Dean of Faculty and Graduate Studies

Unclassified

Security Classification of this page

## REPORT DOCUMENTATION PAGE

1a Report Security Classification UNCLASSIFIED				1b Restrictive Markings			
2a Security Classification Authority				3 Distribution Availability of Report			
2b Declassification/Downgrading Schedule				Approved for public release; distribution is unlimited			
4 Performing Organization Report Number(s) NPS55-90-17				5 Monitoring Organization Report Number(s)			
6a Name of Performing Organization Naval Postgraduate School		6b Office Symbol (If Applicable) OR		7a Name of Monitoring Organization			
6c Address (city, state, and ZIP code) Monterey, CA 93943-5000				7b Address (city, state, and ZIP code)			
8a Name of Funding/Sponsoring Organization Naval Postgraduate School		8b Office Symbol (If Applicable)		9 Procurement Instrument Identification Number 0&MN, Direct Funding			
8c Address (city, state, and ZIP code) Monterey, California  Monterey, CA 93943				10 Source of Funding Numbers			
				Program Element Number	Project No	Task No	Work Unit Accession No
11 Title (Include Security Classification) Modeling and Simulation of Random Shapes by Sculptured Wraparound: Preliminary Report							
12 Personal Author(s) Gaver, D. P. and Jacobs, P. A.							
13a Type of Report Technical		13b Time Covered From To		14 Date of Report (year, month, day) 1990, August		15 Page Count	
16 Supplementary Notation The views expressed in this paper are those of the author and do not reflect the official policy or position of the Department of Defense or the U.S. Government.							
17 Cosati Codes			18 Subject Terms (continue on reverse if necessary and identify by block number)				
Field	Group	Subgroup	Random shapes; Ornstein-Uhlenbeck process; symmetric stable law; Brownian Bridge				
19 Abstract (continue on reverse if necessary and identify by block number)							
Physical objects with boundaries that are irregular—random shapes—occur in profusion in the natural world. Simple mathematical models for random shapes are described. The models are generated by "adding" a random process to a deterministic shape. Ornstein-Uhlenbeck processes are generalized to allow innovation having symmetric stable laws.							
20 Distribution/Availability of Abstract <input checked="" type="checkbox"/> unclassified/unlimited <input type="checkbox"/> same as report <input type="checkbox"/> DTIC users				21 Abstract Security Classification Unclassified			
22a Name of Responsible Individual D. P. Gaver				22b Telephone (Include Area code) (408) 646-2605		22c Office Symbol OR/Gv, OR/Jc	

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted

All other editions are obsolete

security classification of this page

Unclassified

# MODELING AND SIMULATION OF RANDOM SHAPES BY SCULPTURED WRAPAROUND: PRELIMINARY REPORT

Donald P. Gaver

Patricia A. Jacobs

## 1. Introduction

Physical objects with boundaries that are irregular--*random shapes*--occur in profusion in the natural world. Geographical examples range from the continents to small islands, from oceans to lakes to ponds, and from massive ice-covered regions in the Arctic and Antarctic to individual ice floes and icebergs, and their complements, the open spaces or leads in ice fields. There are also important random shapes of biological and medical interest and concern. Those who do remote sensing, either of the atmosphere (clouds), the earth or a planet, from satellites will be interested in describing the shapes observed. Those interested in outlining the shapes of mineral or petroleum deposits encounter also the problem of describing the objects of their search. We propose here some possible alternatives.

It appears that there are relatively few mathematical models for flexibly describing such random shapes. The literature is scattered, and no attempt will be made here to cover it. The writer has examined books by Solomon (1978) and Ripley (1981), but the present focus is different. The work of Kendall and his co-workers may be relevant, cf. Kendall (1985,1989).

Our primary objective at this time is to suggest and explore relatively simple ways of modeling or simulating random shapes that resemble those that occur in nature. Another objective is to describe such shapes as they may

or	
<input checked="checked" type="checkbox"/>	
<input type="checkbox"/>	
on	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



result from an image processing or filtering algorithm's treatment of the basic shape plus a characterization of a noise. Very few detailed analytical results are provided here. Rather, there will be proposals for future work to expand understanding of the basic approaches to be described.

## 2. Random Shapes by Perturbation of Regular Shapes

Many of the two-dimensional irregularly shaped bodies encountered in nature are rather regular or even symmetrical in general outline but possessed of irregular boundaries. We proceed to simulate such figures by starting with a circular disk-like central figure and adding to its perimeter a random process. As will be clear from Figures 1-2, the results are of an interesting qualitative character; that character can be easily changed by manipulating a few parameters. Furthermore, analytical facts can, in various cases, be deduced about features of the shapes so generated. There remains much more to be done in that direction, however.

### Example 1: Shapes with Circular Centrality and Ornstein-Uhlenbeck Boundaries


An especially simple version of our general approach is to represent the boundary of a two-dimensional random shape as follows in polar coordinates:

$$\begin{aligned}\theta(t) &= t, & 0 \leq t \leq 2\pi \\ R(t) &= r + f(X(t))\end{aligned}\tag{1}$$

where  $X(t)$  is an Ornstein-Uhlenbeck (diffusion) process

$$dX(t) = -\alpha X(t) + \sigma dW(t), \quad |\alpha| < 1,\tag{2}$$


with  $\{dW(t)\}$  the increment to a standard Wiener process, and  $f(x) \geq 0$  for any real  $x$ . Useful versions of  $f$  are  $f(x) = e^{kx}$  and  $f(x) = x^2$ . These guarantee that the



**Figure 1.**  $r = 2$ ,  $f(x) = x^2$ ; the parameters of the O.U. process are  $\sigma = 1.0$ ,  $\alpha = 0.4$

perimeter of the random shape is always outside that of the regular shape underlying the process, namely that of a circular disk of radius  $r$ . See Figures 1 and 2 for various realizations of the above; they are of course generated by discretizing time (= angle  $\theta$ ) and replacing  $X(t)$  by an equivalent AR-1 process; the process then starts at  $t \equiv \theta = 0$  with a random draw from a distribution resembling the O.-U. stationary distribution and is plotted going counter-

clockwise around to  $t \equiv \theta = 2\pi$ . The ellipse in each figure is the graph of a circle of radius 2 for the scale of the figure. Note that there is an anticipated (small) jump at the seam at  $t = 0 = 2\pi$  which can be mended in various plausible ways. One such is to force the random boundary function  $X(t)$  to be a *bridge*, so  $X(0) = X(2\pi)$ . This is aesthetically pleasing, but probably unnecessary from a practical viewpoint. It seems natural to call the construction suggested a wraparound process.



**Figure 2.**  $r = 2$ ,  $f(x) = \exp\left(\frac{1}{2}x\right)$ ; the parameters of the O.U. process  
are  $\sigma = 1.0$ ,  $\alpha = 0.4$

Certain analytical results are available directly; others will require more careful attention. For example, let  $A$  be the area of the random shape generated as described. Then, by polar coordinate rules

$$\begin{aligned} A &= \int_0^{2\pi} (r + f(X(t)))^2 \frac{1}{2} dt \\ &= \pi r^2 + r \int_0^{2\pi} f(X(t)) dt + \frac{1}{2} \int_0^{2\pi} f(X(t))^2 dt. \end{aligned} \quad (3)$$

Now take expectations to find

$$E[A] = \pi r^2 + r \int_0^{2\pi} E[f(X(t))] dt + \frac{1}{2} \int_0^{2\pi} E[f(X(t))^2] dt. \quad (4)$$

Suppose  $f(x) = e^{kx}$ ; then if  $X(t)$  is O.U. as in (2) we know that

$$X(t) \sim \mathcal{N}\left(X(0)e^{-\alpha t}, \frac{\sigma^2}{2\alpha}(1 - e^{-2\alpha t})\right) \quad (5)$$

so

$$E\left[e^{+kX(t)}\right] = \exp\left[kX(0)e^{-\alpha t} + \frac{k^2}{2} \cdot \frac{\sigma^2}{2\alpha}(1 - e^{-2\alpha t})\right].$$

Suppose  $X(0) \sim \mathcal{N}\left(0, \frac{\sigma^2}{2\alpha}\right)$ , the stationary distribution. Then, if so, the unconditional distribution of  $X(t) \sim \mathcal{N}\left(0, \frac{\sigma^2}{2\alpha}\right)$  independent of  $t$  for all  $t$ , and we can easily evaluate the mean of the area,  $A$ , for any  $f$ :

$$E[A] = \pi r^2 + 2\pi r \left\{ E[f(X)] + \frac{1}{2} E[f^2(X)] \right\}. \quad (6)$$

For  $f(x) = e^{kx}$  we get



$$E[A] = \pi r^2 + 2\pi r \left\{ e^{k^2 \sigma^2 / 4\alpha} + \frac{1}{2} e^{k^2 \sigma^2 / \alpha} \right\} \quad (7)$$

which shows that as the correlation coefficient  $\alpha$  approaches zero, and the boundaries become exceedingly jagged, the area tends to increase at an exponential rate. Likewise, if  $C$  is a random chord that crosses through the center of the figure then

$$C = 2r + f(X(t)) + f(X(t + \pi)) \quad (8)$$

and if  $X(t)$  is given the stationary distribution then

$$E[C] = 2r + 2E[f(X)], \quad (9)$$

which in the case  $f(x) = e^{kx}$  yields

$$E[C] = 2r + 2e^{k^2 \sigma^2 / 4\alpha} \quad (10)$$

which again suggests the increase in dimension of the shape as  $\alpha$  tends to zero.

Covariances of radii can be obtained analytically, and from these the variance of an area. It is also of interest to ask for the expected length of the maximum central chord.

**Example 2. The previous process, summed.**

There is nothing to prevent one from summing or convex-combining several contributions to the boundary:

$$R(t) = r + \sum f_i(X_i(t)) \quad (11)$$

where, for example,  $X_i(t) \sim \text{O.U.}(\alpha_i, \sigma_i)$  means that the component processes are Ornstein-Uhlenbeck with their individual parameters. Likewise, the correlation and innovation variance parameters can themselves be,

minimally, time-dependent, and possibly even governed by driving stochastic processes, optionally taken to be of the forms

$$\alpha_i(t) = e^{Y_i(t)} / 1 + e^{Y_i(t)} \quad (\text{Logistic}) \quad (12)$$

and

$$\sigma_i(t) = e^{Z_i(t)} \quad (\text{Log - Linear}) \quad (13)$$

with  $Y_i(t)$  and  $Z_i(t)$  themselves taken to be appropriate stochastic processes, not necessarily independent and, most conveniently but not necessarily Gaussian. There are many options here, including that of allowing the radius of the basic circular disk to be itself a random process that, perhaps, tends to grow or shrink in time. A candidate for  $r(t)$  is the radial Brownian motion or Bessel process; see Feller (1966) Chap. X, Section 6, although there are a great many other options. The O.-U. is illustrative and convenient but certainly not intended to be exclusively, if ever, entirely appropriate.

### **Example 3: Perturbing Natural Shapes**

It may be of interest to start with a natural shape, e.g., an ice floe or atmospheric cloud image, and perturb it by a known process, such as the log-O.-U. This could well represent the appearance of the image of the same entity at a different time point, e.g., at a time when a satellite next passes; the difference in apparent shape could be the result of natural processes plus noise.

Suppose a group of such entities exists, each one of which is imaged at time  $t$ , again at  $t+\tau$  at which time shape has been randomly modified as described above, ... and so on with another image take at  $t+k\tau$  with further random perturbation, at each time. Then an important problem is to identify

and track each member of the group over time, under the handicap that shape change can hamper identifiability across different satellite passes. It is proposed to study the above problem, and to investigate the manner in which item identifiability degenerates as the variation in the perturbing boundary process increases.

Note that it is possible to *estimate* the parameters of the perturbing boundary process  $f(X(t))$  by a) first smoothing the figure's perimeter, perhaps by the techniques of Banfield and Raftery (1989), and b) modeling the residual process: the difference of the real/true perimeter and the smoothed perimeter; that difference should be taken orthogonally to the smooth.

**Comment: Relation to "Sculpturing"**

It was Tukey who originally suggested simple ways of lengthening (or shortening) the tails of distributions such as the Gaussian: if  $Z$  is  $\mathcal{N}(0,1)$  then

$$Y = Ze^{hZ}, \quad h > 0 \quad (14)$$

has zero mean but larger variance owing to the extended tail imposed by  $e^{hz}$ . If  $h < 0$  then the tail of the distribution of  $Y$  is shortened. Such an idea could be applied point-wise to  $X(t)$ , e.g., O.-U. from (2) to generate a controllable-shaped boundary: replace  $X(\theta)$  by  $Y(\theta) \equiv Y(t) = X(t)\exp[h \cdot X(t)]$ . One could even allow the sculpturing parameter  $h$  to be angle-dependent, e.g., let  $h(\theta) = -0.2$  for  $0 \leq \theta < \pi$ ,  $h(\theta) = +0.2$  for  $\pi \leq \theta \leq 2\pi$  should result in a figure or shape that is flatter on one side (between 0 and  $\pi$ ) and more jagged on the other (between  $\pi$  and  $2\pi$ ). There is no trouble at all to allow the sculpturing parameter  $h$  to be itself a random process i.e., replace constant  $h$  by  $H(\theta) \equiv H(t)$ . Furthermore, summary statistics are relatively easy to compute if  $X(0)$  is sampled from the stationary distribution of  $\{X(t)\}$ , presuming the latter to exist. Note that the

sculptured figures are exceedingly easy to simulate on a computer: simply generate  $X(t)$ , and then multiply by  $\exp[hX(t)]$  to obtain the boundary value at  $t = \theta$ . [The result is a **sculptured wraparound process** that may be induced to take on natural-appearing shapes by changing a few parameters.]

For an account of general random variable sculpturing see Gaver, et al., (1979) and Gaver (1983). Here we are actually sculpturing a figure or shape, and by a stochastic process realization.

### Application

A report by Banfield and Raftery (1989) suggests application of a non-parametric smoothing technique to better delineate, and identify ice floes. Different satellite images of the same ice floe may differ somewhat, and the B-R procedure should be of help in identifying floes (that have not broken up or experienced edge-alteration) as the same when depicted by different images. Their examples suggest that the procedure works well.

It may be of help to check the effectiveness of B-R smoothing when perturbation or sculpturing of a known principal curve (e.g., circle; or one actually derived from real data) with a boundary or perimeter of known stochastic structure is in place. One simple mechanism for generating such a boundary is to utilize our procedure described above. In order to determine the nature of the parameter for the perturbing process  $f(X(t))$  an examination of residuals (Tukey's "rough") obtained from the estimated principal curve (Tukey's "smooth") should be of use. It is our proposal to attempt such an investigation. It may also be found that procedures less computer-intensive than those of B-R can be nearly as effective.

A general question of applied interest might be the use of remote sensing (by satellite) of ice floes to infer forces causing their (relative) motion, that is, the magnitude of winds or currents between successive satellite passes could account for the changes of relative ice floe positions. Consequently, study of position changes of ice floes might provide a remote inference concerning near-surface forces. For this to work, identifiability of individual ice floes seems of considerable importance. Such an approach, to infer surface wind speed from satellite sensed white-cap cover, has been taken by Monahan and O'Muircheartaigh (1980).

### References and Bibliography

J. D. Banfield and A. E. Raftery. "Ice floe identification in satellite images using mathematical morphology and clustering about principal curves." Technical Report 172, Department of Statistics, GN22, University of Washington, August 1989.

D. P. Gaver and M. Acar. "Analytic hazard representations for use in reliability, mortality, and simulation studies. *Commun. Statist. Simula. Computa.* B8 (2), 1979, pp. 91-111.

W. Feller. *An Introduction to Probability Theory and its Applications*, Vol. II, John Wiley, New York: 1966.

D. P. Gaver. "Stochastic modeling: ideas and techniques" in *Probability Theory and Computer Science*, ed. G. Louchard and G. Latouche. Academic Press, New York, 1983.

D. G. Kendall. "A survey of the statistical theory of shape," (with discussion) *Statistical Science*, 4, (2), 1989, pp. 87-120.

D. G. Kendall. "Exact distributions for shapes of random triangles in convex sets." *Adv. Appl. Prob.*, 17, pp. 308-329.

E. C. Monahan and I. G. O'Muircheartaigh. "Optimal power-law description of oceanic whitecap coverage dependence on wind speed." *J. Phys. Oceanography*, 10, pp. 2084-2099.

B. D. Ripley. *Spatial Statistics*, Wiley, New York, 1981.

H. Solomon. *Geometric Probability*, Society for Industrial and Applied Mathematics, Philadelphia, 1978.

## APPENDIX A. GENERALIZING THE ORNSTEIN-UHLENBECK PROCESS

The purpose of this note is to examine a model analogous to the Ornstein-Uhlenbeck (O.-U.) process, in which innovations are allowed to have *symmetric stable laws*, cf. Feller (1968), rather than be simply Gaussianly distributed bits of standard Wiener processes. In particular, such processes can have Cauchy innovations. The purpose of this is to introduce processes that give rise to long tails that can provide more adjustable behavior for image boundaries.

Thus for arbitrarily small  $h > 0$  we write

$$X(t+h) = X(t) - \rho h X(t) + \tau h \epsilon_t \quad (A-1)$$

where  $\epsilon_t$  has, concretely, the Cauchy density

$$\epsilon_t \sim \frac{1}{\pi} \left( \frac{1}{1+x^2} \right). \quad (A-2)$$

For this distribution the characteristic function (ch. fcn.) is

$$E[e^{i\theta\epsilon_t}] = e^{-|\theta|}. \quad (A-3)$$

In the most general case, that of a symmetric stable law,

$$E[e^{i\theta\epsilon_t}] = e^{-|\theta|^\alpha}, 1 \leq \alpha \leq 2; \quad (A-4)$$

if  $\alpha=2$  it is recognized that innovations are Gaussian.

Put

$$\varphi(\theta, t) = E[e^{i\theta X(t)}] \quad (A-5)$$

the ch. fcn. of  $X(t)$ ; we keep implicit the initial condition  $X(0)$ . In general  $\theta$  must be a real number in order that the rhs of (A-4) be finite. Derive an equation for  $\varphi$  as follows. Write

$$\begin{aligned} E\left[e^{i\theta X(t+h)}\right] &= E\left[e^{i\theta\{X(t)(1-\rho h)+\tau h\epsilon_t\}}\right] \\ &= E\left[e^{i\theta\{X(t)(1-\rho h)\}}\right] E\left[e^{i\theta\tau h\epsilon_t}\right] = \varphi(\theta(1-\rho h), t) e^{-h\xi|\theta|^\alpha} \quad (A-6) \end{aligned}$$

by the assumed independence of successive innovations; note that it is necessary to scale innovations so that  $\tau^\alpha h^\alpha = \xi h$  for fixed  $\xi$ , or  $\tau = \xi^{1/\alpha} h^{(1-\alpha)/\alpha}$  for fixed  $\xi$ . If  $h$  becomes small,

$$E\left[e^{i\theta X(t+h)}\right] \equiv \varphi(\theta, t+h) \approx \left[\varphi(\theta, t) + \left(\frac{\partial}{\partial \theta} \varphi(\theta, t)\right)(-\rho\theta)h + O(h^2)\right] \left[1 - h\xi|\theta|^\alpha + O(h^2)\right] \quad (A-7)$$

This done, then

$$\frac{\varphi(\theta, t+h) - \varphi(\theta, t)}{h} = -\rho\theta \frac{\partial \varphi}{\partial \theta} - \xi|\theta|^\alpha \varphi + O(h) \quad (A-8)$$

and if  $h \rightarrow 0$  the following quasi-linear first-order partial differential equation results:

$$\frac{\partial \varphi}{\partial t} + \rho\theta \frac{\partial \varphi}{\partial \theta} = -\xi|\theta|^\alpha \varphi. \quad (A-9)$$

### Steady-State or Long-Run Solution

Suppose

$$\lim_{t \rightarrow \infty} \varphi(\theta, t) = \varphi^*(\theta) \quad (A-10)$$

is a bona-fide ch. fcn. of an honest distribution. This is at least plausible since  $\rho > 0$  and there is a restoring force tendency in the dynamics of (A-1). We then have the ordinary first-order differential equation



$$\rho\theta \frac{d\varphi^\#}{d\theta} = -\xi|\theta|^\alpha \varphi^\# \quad (A-11)$$

or

$$\frac{d\varphi^\#}{\varphi^\#} = -\frac{\xi}{\rho} \frac{|\theta|}{\theta} |\theta|^{\alpha-1} d\theta = -\frac{\xi}{\rho} s(\theta) |\theta|^{\alpha-1} d\theta \quad (A-12)$$

where

$$s(\theta) = \begin{cases} +1 & \text{if } \theta \geq 0 \\ -1 & \text{if } \theta < 0 \end{cases} \quad (A-13)$$

Write (A-12) as

$$\frac{d\varphi^\#}{\varphi^\#} = -\frac{\xi}{\rho} |\theta|^{\alpha-1} d|\theta|$$

and integrate to obtain

$$\varphi^\#(\theta) = \exp\left[-\frac{\xi}{\rho\alpha} |\theta|^\alpha\right] \quad (A-14)$$

This is the ch.fcn. of a symmetric stable law, as before. If  $\alpha=2$ ,  $\varphi^\#$  is the ch. fcn. of a Gaussian/Normal with variance  $\xi/\rho$ , as is used earlier, (2). If  $\alpha=1$ ,  $\varphi^\#$  represents a Cauchy distribution with scale parameter  $\xi/\rho$ . Other special cases may exist, but are far less familiar and convenient.

### Transient Solution

The equation (A-9) can be solved by the standard method of characteristics; see Street (1973). Another approach that reveals the underlying structure of the process is this: express (A-1) as

$$dX(t) = -\rho X(t)dt + \tau \epsilon_t dt \quad (A-15)$$

or formally as

$$\frac{dX}{dt} = -\rho X(t) + \tau \epsilon_t. \quad (A-16)$$

Note that we can allow  $\rho$  to depend on  $t$  at no cost:

$$\frac{dX}{dt} = -\rho(t)X(t) + \tau \epsilon_t. \quad (A-17)$$

Use the standard integrating factor:

$$\frac{d}{dt} \left[ X(t) e^{\int_0^t \rho(x) dx} \right] = \tau \epsilon_t e^{\int_0^t \rho(x) dx}; \quad (A-18)$$

so if  $X(0) = X_0$ ,

$$X(t) = X_0 e^{\int_0^t \rho(x) dx} + \int_0^t e^{-\int_y^t \rho(x) dx} \tau \epsilon_y dy. \quad (A-19)$$

Clearly  $X(t)$  has the structure of a weighted sum of components; a typical member of the integrand being

$$e^{-\int_y^t \rho(x) dx} \tau \epsilon_y dy. \quad (A-20)$$

Therefore

$$\begin{aligned} E \left[ e^{i\theta X(t)} \right] &= E \left[ \exp \left\{ i\theta \left[ X_0 e^{\int_0^t \rho(x) dx} + \int_0^t e^{-\int_y^t \rho(x) dx} \tau \epsilon_y dy \right] \right\} \right] \\ &= E \left[ \exp \left\{ i\theta X_0 e^{\int_0^t \rho(x) dx} \right\} \right] \cdot E \left[ \exp \left\{ i\theta \int_0^t e^{-\int_y^t \rho(x) dx} \tau \epsilon_y dy \right\} \right] \end{aligned} \quad (A-21)$$

Now think of the integral  $\int_0^t e^{-\int_y^t \rho(x)dx} \epsilon_y dy$  as approximated by a sum of elements, a typical value being (A-20). Use the independence of successive  $\epsilon_y$  contributions to obtain

$$E \left[ \exp \left\{ i \theta \int_0^t e^{-\int_y^t \rho(x)dx} \epsilon_y dy \right\} \right] = \exp \left\{ -|\theta|^\alpha \int_0^t e^{-\alpha \int_y^t \rho(x)dx} \xi dy \right\} \quad (A-22)$$

for, by the fact that  $\epsilon_y$  obeys a stable law

$$E \left[ \exp \left\{ i \theta e^{-\int_y^t \rho(x)dx} \epsilon_y dy \right\} \right] = \exp \left\{ -|\theta|^\alpha e^{-\alpha \int_y^t \rho(x)dx} \tau^\alpha (dy)^\alpha \right\} \quad (A-23)$$

again we have put  $\xi dy = \tau^\alpha (dy)^\alpha$  just as was done before. It follows that the solution to (A-9) must be

$$\varphi(\theta, t) = \exp \left\{ i \theta X_0 e^{-\rho t} - |\theta|^\alpha \frac{\xi}{\alpha \rho} (1 - e^{-\alpha \rho t}) \right\} \quad (A-24)$$

Thus for fixed  $t$ ,

$$X(t) \sim \text{Stable} \left( \text{center} = X_0 e^{-\rho t}, \text{scale} = \left( \frac{\xi}{\alpha \rho} (1 - e^{-\alpha \rho t}) \right)^{\frac{1}{\alpha}}; \text{index} = \alpha \right); \quad (A-25)$$

the center is the mean and the scale is the standard deviation in the Gauss/Normal case  $\alpha = 2$ . Of course (A-25) agrees with (A-14) as  $t \rightarrow \infty$ , and differentiation will show that (A-24) solves (A-9). *Note:* the rate of approach to steady state,  $\alpha \rho$ , increases with  $\alpha$ , so the Gauss/Normal case (classical O.-U.) approaches steady-state more quickly than does the analogous Cauchy-driven O.-U. with same  $\rho$ .

## Simulation

In order to simulate  $X(t)$  it will be necessary to discretize time into, say,  $h$ -increments; the result will be a path or trajectory  $X_h(t)$ . In order to obtain the desired marginal distributions, e.g., (A-25) simply use (A-1), i.e.

$$X(t+h) = X(t) - \rho h X(t) + \tau \epsilon_t, \quad t = 0, h, 2h, \dots$$

where  $\tau = \xi^{1/\alpha} h^{(1-\alpha)/\alpha}$  and  $\epsilon_t$  is an  $\alpha$ -stable random variable with unit scale.

For example(s):

$\alpha = 2$ ;  $\epsilon_t$  is a draw from  $e^{-0.5x^2} / \sqrt{2\pi}$  (Normal density)

$\alpha = 1$ ;  $\epsilon_t$  is a drawn from  $\left[ \pi(1+x^2) \right]^{-1}$  (Cauchy density).

For fixed  $\xi$  and  $\rho$  the paths of the Cauchy O.-U. should tend to exhibit more long-range order than those of the Gauss O.-U. The option of choosing  $\rho = \rho(t)$  allows further freedom in adjusting path behavior.

## Reference

R. L. Street (1973) *Analysis and Solution of Partial Differential Equations*.  
Brooks-Cole Publishing Co. (Wadsworth), Monterey, CA.

## APPENDIX B. BRIDGE PROCESSES

One feature common to Figures 1 and 2 is the jump that appears at the seam of  $t=0=2\pi$ . The purpose of this Appendix is to describe one procedure to obtain shapes that are "seamless." To this end processes for the Ornstein-Uhlenbeck (O.-U.) and Cauchy autoregressive processes are introduced that correspond to the Brownian Bridge.

Let  $\{W(t); 0 \leq t \leq 1\}$  be a standard Brownian motion process with  $W(0)=0$ . Recall that a Brownian Bridge can be obtained by the transformation

$$B(t) = W(t) - tW(1), \quad 0 \leq t \leq 1;$$

note that  $B(1) = B(0) = 0$ ; cf. Karlin and Taylor (1980).

In a similar spirit let  $\{X(t), t \geq 0\}$  be an O.-U. process or a generalization of it as described in Appendix A with  $X(0) = 0$ . We will define a Bridge process as

$$U(t) = X(t) - tX(1), \text{ for } 0 \leq t \leq 1.$$

Note that  $X(0) = X(1) = 0$ . One way not to have  $X(0) = X(1) = 0$  is to add a random variable  $U^0$  to  $U(t)$  having the stationary distribution of  $\{X(t); t \geq 0\}$ .

Let  $\{X(t); t \geq 0\}$  be an O.-U. process with drift coefficient  $-\rho$ , infinitesimal variance  $\sigma^2$ , and  $X(0) = 0$ . Set

$$Y(t) = X(t) - tX(1), \quad 0 \leq t \leq 1.$$

$\{Y(t); t \geq 0\}$  is a Gaussian process with mean 0. Further, for  $s < t$

$$\begin{aligned} E[Y(t)Y(s)] &= E[(X(t) - tX(1))(X(s) - sX(1))] \\ &= E[X(t)X(s)] - tE[X(1)X(s)] - sE[X(t)X(1)] + tsE[X(1)^2] \\ &= \frac{\sigma^2}{2\rho} \left\{ e^{-\rho(t+s)} [e^{2\rho s} - 1] - te^{-\rho(1+s)} [e^{2\rho s} - 1] - se^{-\rho(1+t)} [e^{2\rho t} - 1] + tse^{-2\rho} [e^{2\rho} - 1] \right\}. \end{aligned}$$

Thus,

$$\begin{aligned} E[Y(t)^2] &= \text{Var}[Y(t)] \\ &= \frac{\sigma^2}{2\rho} \left[ (1 - e^{-2\rho t}) - 2te^{-\rho(1+t)}(e^{2\rho t} - 1) + t^2(1 - e^{-2\rho}) \right]. \end{aligned}$$

### Simulation.

The O.-U. Bridge can be simulated as follows.  $X^\circ$  is generated from a normal distribution having mean 0 and variance  $\sigma^2/2\rho$ . The Ornstein-Uhlenbeck process is simulated as  $X(0)=0$

$$X(t+h) = X(t) - \rho h X(t) + \sqrt{h} \epsilon_t$$

where  $\{\epsilon_t\}$  is a sequence of independent normal random variables with mean 0 and variance  $\sigma^2$ . A bridge is created by calculating

$$\tilde{U}(t) = X(t) - tX(1) \quad 0 \leq t \leq 1.$$

The random variable  $X^\circ$  is added to  $\tilde{U}(t)$  to obtain the process

$$U(t) = \tilde{U}(t) + X^\circ.$$

In the figure 1B, the radius of the shape at time  $2\pi t$  is taken to be

$$R + |U(t)|.$$

The parameters of the simulation are  $R = 2$ ,  $\rho = 0.4$ ,  $\sigma = 1$  and  $h = .005$ . The ellipse in the center of the figure is the circle of radius 2 for the scale of the picture.

The Cauchy Bridge is simulated as follows.  $X^\circ$  is generated from a Cauchy distribution having density function

$$f^\circ(y) = (\xi / \rho)^{-1} \frac{1}{\pi} \left[ 1 + (y\rho / \xi)^2 \right]^{-1}.$$

In the simulation,

$$\begin{aligned} X(0) &= 0 \\ X(t+h) &= X(t) - \rho h X(t) + \xi \epsilon(t) h \quad 0 \leq t \leq 1 \end{aligned}$$

where  $\{\epsilon(t)\}$  is a sequence of independent standard Cauchy random variables.

A Bridge is created by calculating

$$\tilde{U}(t) = X(t) - tX(1). \quad 0 \leq t \leq 1.$$

Finally, a Bridge not tied down at 0 is obtained by setting

$$U(t) = \tilde{U}(t) + X^\circ.$$

In Figure 2B the radius of the shape at time  $2\pi t$  is taken to be

$$R + |U(t)|.$$

The parameters of the simulation are  $R = 2$ ,  $\rho = 0.4$ ,  $h = 0.005$ , and  $\xi = 0.8$ . Once again the ellipse in the center is the circle of radius 2 for the scale of the figure.

## Reference

S. Karlin and H. M. Taylor (1981) *A Second Course in Stochastic Processes*. Academic Press Inc., New York.

$$|O-U \text{ BRIDGE}| + RV + R$$

$$R=2; \text{RHO}=0.4; H=.005; \text{SIG}=1$$




Figure 1B



|CAUCHY BRIDGE|+RV+R

R=2;RHO=0.4;H=.005;PSI=.8




Figure 2B

## INITIAL DISTRIBUTION LIST

1. Library (Code 0142).....2  
Naval Postgraduate School  
Monterey, CA 93943-5000
2. Defense Technical Information Center.....2  
Cameron Station  
Alexandria, VA 22314
3. Office of Research Administration (Code 012) .....1  
Naval Postgraduate School  
Monterey, CA 93943-5000
4. Prof. Peter Purdue.....1  
Code OR-Pd  
Naval Postgraduate School  
Monterey, CA 93943-5000
5. Department of Operations Research (Code 55).....1  
Naval Postgraduate School  
Monterey, CA 93943-5000
6. Prof. Donald Gaver, Code OR-Gv.....15  
Naval Postgraduate School  
Monterey, CA 93943-5000
7. Prof. Patricia Jacobs .....15  
Code OR/Jc  
Naval Postgraduate School  
Monterey, CA 93943-5000
8. Center for Naval Analyses.....1  
4401 Ford Avenue  
Alexandria, VA 22302-0268
9. Dr. David Brillinger.....1  
Statistics Department  
University of California  
Berkeley, CA 94720

10. Dr. R. Gnanadesikan .....1  
Bellcore  
435 South Street  
Morris Township NJ 07960
11. Prof. Bernard Harris .....1  
Dept. of Statistics  
University of Wisconsin  
610 Walnut Street  
Madison, WI 53706
12. Prof. W. M. Hinich .....1  
University of Texas  
Austin, TX 78712
13. Prof. I. R. Savage.....1  
Dept. of Statistics  
Yale University  
New Haven, CT 06520
14. Prof. W. R. Schucany.....1  
Dept. of Statistics  
Southern Methodist University  
Dallas, TX 75222
15. Prof. D. C. Siegmund.....1  
Dept. of Statistics  
Sequoia Hall  
Stanford University  
Stanford, CA 94305
16. Prof. H. Solomon .....1  
Department of Statistics  
Sequoia Hall  
Stanford University  
Stanford, CA 94305
17. Dr. Ed Wegman.....1  
George Mason University  
Fairfax, VA 22030
18. Dr. P. Welch .....1  
IBM Research Laboratory  
Yorktown Heights, NY 10598

19. Dr. Neil Gerr .....1  
Office of Naval Research  
Arlington, VA 22217
20. Prof. Roy Welsch.....1  
Sloan School  
M.I.T.  
Cambridge, MA 02139
21. Dr. J. Abrahams .....1  
Code 1111, Room 607  
Mathematical Sciences Division  
Office of Naval Research  
800 North Quincy Street  
Arlington, VA 22217-5000
22. Prof. J. R. Thompson.....1  
Dept. of Mathematical Science  
Rice University  
Houston, TX 77001
23. Dr. P. Heidelberger.....1  
IBM Research Laboratory  
Yorktown Heights  
New York, NY 10598
24. Prof. M. Leadbetter.....1  
Department of Statistics  
University of North Carolina  
Chapel Hill, NC 27514
25. Prof. D. L. Iglehart.....1  
Dept. of Operations Research  
Stanford University  
Stanford, CA 94350
26. Prof. J. B. Kadane.....1  
Dept. of Statistics  
Carnegie-Mellon University  
Pittsburgh, PA 15213

27. Prof. J. Lehoczky .....1  
Department of Statistics  
Carnegie-Mellon University  
Pittsburgh, PA 15213
28. Dr. J. Maar (R513) .....1  
National Security Agency  
Fort Meade, MD 20755-6000
29. Prof. M. Mazumdar .....1  
Dept. of Industrial Engineering  
University of Pittsburgh  
Pittsburgh, PA 15235
30. Prof. M. Rosenblatt .....1  
Department of Mathematics  
University of California, San Diego  
La Jolla, CA 92093
31. Prof. H. Chernoff .....1  
Department of Statistics  
Harvard University  
1 Oxford Street  
Cambridge, MA 02138
32. Dr. T. J. Ott .....1  
Bellcore  
435 South Street  
Morris Township, NJ 07960
33. Dr. Alan Weiss .....1  
AT&T Bell Laboratories  
Mountain Avenue  
Murray Hill, NJ 07974
34. Prof. Joseph R. Gani .....1  
Mathematics Department  
University of California  
Santa Barbara, CA 93106
35. Prof. Frank Samaniego .....1  
Statistics Department  
University of California  
Davis, CA 95616

36. Dr. James McKenna.....1  
AT&T Bell Laboratories  
Mountain Avenue  
Murray Hill, NJ 07974
37. Commander.....1  
Attn: G. F. Kramer  
Naval Ocean Systems Center  
Code 421  
San Diego, CA 92152-5000
38. Prof. Tom A. Louis.....1  
School of Public Health  
University of Minnesota  
Mayo Bldg. A460  
Minneapolis, MN 55455
39. Dr. Nan Laird.....1  
Biostatistics Dept.  
Harvard School of Public Health  
677 Huntington Ave.  
Boston, MA 02115
40. Dr. Marvin Zelen.....1  
Biostatistics Department  
Harvard School of Public Health  
677 Huntington Ave.  
Boston, MA 02115
41. Dr. John Orav.....1  
Biostatistics Department  
Harvard School of Public Health  
677 Huntington Ave.  
Boston, MA 02115
42. Prof. R. Douglas Martin.....1  
Department of Statistics, GN-22  
University of Washington  
Seattle, WA 98195

43. Prof. W. Stuetzle .....1  
Department of Statistics  
University of Washington  
Seattle, WA 98195
44. Prof. F. W. Mosteller .....1  
Department of Statistics  
Harvard University  
1 Oxford St.  
Cambridge, MA 02138
45. Dr. D. C. Hoaglin .....1  
Department of Statistics  
Harvard University  
1 Oxford Street  
Cambridge, MA 02138
46. Prof. N. D. Singpurwalla .....1  
George Washington University  
Washington, DC 20052
47. Center for Naval Analysis .....1  
2000 Beauregard Street  
Alexandria, VA 22311
48. Prof. George S. Fishman .....1  
Curr. in OR & Systems Analysis  
University of North Carolina  
Chapel Hill, NC 20742
49. Dr. Alan F. Petty .....1  
Code 7930  
Naval Research Laboratory  
Washington, DC 20375
50. Prof. Bradley Efron .....1  
Statistics Dept.  
Sequoia Hall  
Stanford University  
Stanford, CA 94305

51. Prof. Carl N. Morris.....1  
Statistics Department  
Harvard University  
1 Oxford St.  
Cambridge, MA 02138
52. Dr. John E. Rolph.....1  
RAND Corporation  
1700 Main St.  
Santa Monica, CA 90406
53. Prof. Linda V. Green.....1  
Graduate School of Business  
Columbia University  
New York, NY 10027
54. Dr. David Burman.....1  
AT&T Bell Telephone Laboratories  
Mountain Avenue  
Murray Hill, NJ 07974
55. Dr. Ed Coffman.....1  
AT&T Bell Telephone Laboratories  
Mountain Avenue  
Murray Hill, NJ 07974
56. Prof. William Jewell.....1  
Operations Research Department  
University of California, Berkeley  
Berkeley, CA 94720
57. Prof. D. C. Siegmund.....1  
Dept. of Statistics  
Sequoia Hall  
Stanford University  
Stanford, CA 94305
58. Operations Research Center, Rm E40-164.....1  
Massachusetts Institute of Technology  
Attn: R. C. Larson and J. F. Shapiro  
Cambridge, MA 02139



59. Arthur P. Hurter, Jr. ....1  
Professor and Chairman  
Dept. of Industrial Engineering and Management Sciences  
Northwestern University  
Evanston, IL 60201-9990
60. Institute for Defense Analysis.....1  
1800 North Beauregard  
Alexandria, VA 22311
61. Prof. J. W. Tukey .....1  
Statistics Dept., Fine Hall  
Princeton University  
Princeton, NJ 08540
62. Dr. Daniel H. Wagner .....1  
Station Square One  
Paoli, PA 19301
63. Dr. Colin Mallows.....1  
AT&T Bell Telephone Laboratories  
Mountain Avenue  
Murray Hill, NJ 07974
64. Dr. D. Pregibon.....1  
AT&T Bell Telephone Laboratories  
Mountain Avenue  
Murray Hill, NJ 07974
65. Dr. Jon Kettenring.....1  
Bellcore  
435 South Street  
Morris Township, NJ 07960
66. Prof. David L. Wallace .....1  
Statistics Dept.  
University of Chicago  
5734 S. University Ave.  
Chicago, IL 60637
67. Dr. S. R. Dalal.....1  
AT&T Bell Telephone Laboratories  
Mountain Avenue  
Murray Hill, NJ 07974

68. Dr. M. J. Fischer .....1  
Defense Communications Agency  
1860 Wiehle Avenue  
Reston, VA 22070
69. Dr. Prabha Kumar .....1  
Defense Communications Agency.....1  
1860 Wiehle Avenue  
Reston, VA 22070
70. Dr. B. Doshi .....1  
AT&T Bell Laboratories  
HO 3M-335  
Holmdel, NJ 07733
71. Dr. D. M. Lucantoni.....1  
AT&T Bell Laboratories  
Holdmel, NJ 07733
72. Dr. V. Ramaswami .....1  
MRE 2Q-358  
Bell Communications Research, Inc.  
435 South Street  
Morristown, NJ 07960
73. Prof. G. Shantikumar .....1  
The Management Science Group  
School of Business Administration  
University of California  
Berkeley, CA 94720
74. Dr. D. F. Daley .....1  
Statistic Dept. (I.A.S.)  
Australian National University  
Canberra, A.C.T. 2606  
AUSTRALIA
75. Dr. Guy Fayolle.....1  
I.N.R.I.A.  
Dom de Voluceau-Rocquencourt  
78150 Le Chesnay Cedex  
FRANCE

76. Professor H. G. Daellenbach.....1  
Department of Operations Research  
University of Canterbury  
Christchurch, NEW ZEALAND
77. Koh Peng Kong.....1  
OA Branch, DSO  
Ministry of Defense  
Blk 29 Middlesex Road  
SINGAPORE 1024
78. Professor Sir David Cox.....1  
Nuffield College  
Oxford, OXI INF  
ENGLAND
79. Dr. A. J. Lawrence.....1  
Department of Mathematical Statistics  
University of Birmingham  
P. O. Box 363  
Birmingham B15 2TT  
ENGLAND
80. Dr. F. P. Kelly.....1  
Statistics Laboratory  
16 Mill Lane  
Cambridge  
ENGLAND
81. Dr. R. J. Gibbens.....1  
Statistics Laboratory  
16 Mill Lane  
Cambridge  
ENGLAND
82. Dr. John Copas.....1  
Dept. of Mathematical Statistics  
University of Birmingham  
P. O. Box 363  
Birmingham B15 2TT  
ENGLAND

83. Dr. D. Vere-Jones.....1  
Dept. of Mathematics  
Victoria University of Wellington  
P. O. Box 196  
Wellington  
NEW ZEALAND
84. Prof. Guy Latouche.....1  
University Libre Bruxelles  
C. P. 212  
Blvd. De Triomphe  
B-1050 Bruxelles  
BELGIUM